

# MASS BOUNDS FOR MULTIDIMENSIONAL CHARGED DILATONIC BLACK HOLES

U. BLEYER<sup>1</sup> and V. D. IVASHCHUK<sup>2</sup>

WIP-Gravitationsprojekt, Universität Potsdam  
An der Sternwarte 16  
D-14482 Potsdam, Germany

## Abstract

The multidimensional charged dilatonic black hole solution with  $n$  internal Ricci-flat spaces is considered. The bound on the mass of the black hole is obtained. In the strong dilatonic coupling limit the critical mass becomes zero. The case  $n = \infty$  is also considered.

---

<sup>1</sup>E-mail address: UBleyer@aip.de. This work was supported by WIP grant 016659.

<sup>2</sup>Permanent address: Center for Surface and Vacuum Research, 8 Kravchenko str., Moscow, 117331, Russia, e-mail: mel@cvsi.uucp.free.msk.su. This work was supported in part by DFG grant 436 RUS - 113-7-2.

# 1 Introduction

In [1] Myers and Perry obtained the multidimensional  $O(d+1)$ -symmetric analogue of the well-known Reissner-Nordström charged black hole solution. In [2,3] the generalization of the Myers-Perry solution to the case of  $n$  Ricci-flat internal spaces was obtained. Special cases of the solution [2,3] were considered earlier in the following publications: [4,5] ( $d = 2$ ,  $\lambda = 0$ ), [6] ( $d = 2$ ,  $n = 1$ ), [7] ( $d \geq 2$ ,  $\lambda = 0$ ). (In [8] the special case of solution [2,3] with zero electric and scalar charges was obtained.)

The model considered in [2,3] is described by the action

$$S = \int d^D x \sqrt{|g|} \left\{ \frac{1}{2\kappa^2} R[g] - \frac{1}{2\kappa^2} \partial_M \varphi \partial_N \varphi g^{MN} - \frac{1}{4} \exp(2\lambda\varphi) F_{MN} F^{MN} \right\}, \quad (1.1)$$

where  $g = g_{MN} dx^M \otimes dx^N$  is the metric,  $F = \frac{1}{2} F_{MN} dx^M \wedge dx^N = dA$  is the strength of the electromagnetic field and  $\varphi$  is the scalar field (dilaton field). Here  $\lambda$  is the dilaton coupling constant and  $\kappa$  denotes Einstein's gravitational constant. The action (1.1) describes for certain values of the coupling constant  $\lambda$  and space-time dimension  $D$  a lot of interesting physical models including standard Kaluza-Klein theory, dimensionally reduced Einstein-Maxwell theory, certain sectors of supergravity theories etc. For

$$\lambda^2 = \frac{1}{2}, \quad D = 10, \quad (1.2)$$

the action (1.1) describes a part of the bosonic sector for the  $N = 1$  ten-dimensional Einstein-Yang-Mills supergravity that occurs in the low energy limit of superstring theory [9].

In this paper we consider as special case of the solution [2,3] a charged dilaton black hole with  $n$  internal Ricci-flat spaces. Objects of such sort are very popular in the literature (see for example [2,6,10-14]). Here we present bounds on the mass of this black hole. For  $D = 4$ ,  $d = 2$  such a bound was obtained by Gibbons and Wells in [14]. We also consider an interesting special

case of the charged black hole solution with infinite number of internal spaces  $n = \infty$ . We note that to our knowledge the Einstein equations in the infinite-dimensional space were considered first by Kalitzin [15]. Infinite-dimensional spherically-symmetric and cosmological solutions were also presented in [3] and [16] respectively.

## 2 Spherically Symmetric Solutions

The field equations, corresponding to the action (1.1), have the following form

$$R_{MN} - \frac{1}{2}g_{MN}R = \kappa^2 T_{MN}, \quad (2.1)$$

$$\square\varphi - \frac{\kappa^2}{2}\lambda \exp(2\lambda\varphi)F_{MN}F^{MN} = 0, \quad (2.2)$$

$$\nabla_M(\exp(2\lambda\varphi)F^{MN}) = 0, \quad (2.3)$$

where

$$\begin{aligned} T_{MN} = & \frac{1}{\kappa^2}(\partial_M\varphi\partial_N\varphi - \frac{1}{2}g_{MN}\partial_P\varphi\partial^P\varphi) \\ & + \exp(2\lambda\varphi)(F_{MP}F_N{}^P - \frac{1}{4}g_{MN}F_{PQ}F^{PQ}). \end{aligned} \quad (2.4)$$

Here we consider the spherically  $O(d+1)$ -symmetric solutions of the field equations (2.1)-(2.3) obtained in [2,3]. In this paper the notations of ref. [3] are used. The solution [4] is defined on the manifold

$$M = M^{(2+d)} \times M_1 \times \dots \times M_n, \quad (2.5)$$

and has the following form

$$\begin{aligned} g = & -f_1^{(D-3)/A(\lambda)} f_\varphi^{2\lambda} d\bar{t} \otimes d\bar{t} \\ & + f_1^{-1/A(\lambda)} (f_2^{-1} f_\varphi^{2\lambda} f^2)^{1/(1-d)} [f_2 du \otimes du + d\Omega_d^2] \\ & + \sum_{i=1}^n f_1^{-1/A(\lambda)} \exp(2A_i u + 2D_i) g^{(i)}, \end{aligned} \quad (2.6)$$

$$F = Qf_1 du \wedge d\bar{t}, \quad (2.7)$$

$$\exp \varphi = f_1^{(2-D)\lambda/2A(\lambda)} f_\varphi. \quad (2.8)$$

where  $M^{(2+d)}$  is a  $(2+d)$ -dimensional space-time ( $d \geq 2$ ), the  $(M_i, g^{(i)})$  are Ricci-flat manifolds ( $g^{(i)}$  is the metric on  $M_i$ ),  $\dim M_i = N_i$  ( $i = 1, \dots, n$ ),  $d\Omega_d^2$  is the canonical metric on the  $d$ -dimensional sphere  $S^d$ . In (2.6)-(2.8)

$$f_1 = f_1(u) = C_1(D-2)/\kappa^2 Q^2 A(\lambda) \sinh^2(\sqrt{C_1}(u - u_1)), \quad (2.9)$$

$$f_2 = f_2(u) = C_2/(d-1)^2 \sinh^2(\sqrt{C_2}(u - u_2)), \quad (2.10)$$

$$f_\varphi = f_\varphi(u) = \exp(Bu + D_\varphi), \quad (2.11)$$

$$f = f(u) = \exp\left[\sum_{i=1}^n N_i(A_i u + D_i)\right], \quad (2.12)$$

$$A(\lambda) = D - 3 + \lambda^2(D - 2), \quad (2.13)$$

and  $Q \neq 0$ ,  $D_i$ ,  $D_\varphi$ ,  $u_1$ ,  $u_2$  are constants and the parameters  $C_1$ ,  $C_2$ ,  $B$ ,  $A_i$  satisfy the relation

$$\begin{aligned} \frac{C_2 d}{d-1} = & \frac{C_1(D-2)}{D-3+\lambda^2(D-2)} + B^2(1+\lambda^2) \\ & + \frac{1}{d-1}(\lambda B + \sum_{i=1}^n A_i N_i)^2 + \sum_{i=1}^n A_i^2 N_i. \end{aligned} \quad (2.14)$$

### 3 Dilatonic Charged Black Hole

Now, we are looking for a black hole solution, that means for the special case of a configuration with external horizon. To do so, we take the solution (2.6)-(2.14) with the parameters

$$C_1 = C_2 = C > 0, \quad u_2 = 0, \quad u_1 = -u_0 < 0, \quad (3.1)$$

$$A_i/\sqrt{C} = -1/A(\lambda), \quad B/\sqrt{C} = -\lambda(D-2)/A(\lambda). \quad (3.2)$$

Introducing the parameters

$$B_{\pm} = \frac{\kappa|Q|\sqrt{A(\lambda)}}{(d-1)\sqrt{D-2}} \exp(\pm\sqrt{C}u_0) \quad (3.3)$$

and reparametrizing the time and radial coordinates

$$\bar{t} = \frac{\kappa|Q|\sqrt{A(\lambda)} \sinh(\sqrt{C}u_0)}{\sqrt{C(D-2)}} t, \quad (3.4)$$

$$r^{d-1} = \frac{\kappa|Q|\sqrt{A(\lambda)} \sinh(\sqrt{C}(u+u_0))}{(d-1)\sqrt{(D-2)} \sinh(\sqrt{C}u)}, \quad (3.5)$$

we get the following formulas for the solution ( $\lambda \neq 0$ )

$$g = -f_+ f_-^{1+2\alpha_t} dt \otimes dt + f_-^{2\alpha_r} \left[ \frac{dr \otimes dr}{f_+ f_-} + r^2 d\Omega_d^2 \right] + f_-^{-2/A(\lambda)} \sum_{i=1}^n g^{(i)}, \quad (3.6)$$

$$F = Qr^{-d} dt \wedge dr, \quad (3.7)$$

$$\exp(2\lambda\varphi) = f_-^{2\alpha_t}. \quad (3.8)$$

Here

$$f_{\pm} = f_{\pm}(r) = 1 - \frac{B_{\pm}}{r^{d-1}}, \quad (3.9)$$

$$\alpha_t = -\lambda^2(D-2)/A(\lambda), \quad \alpha_r = \frac{1}{d-1} - \frac{1}{A(\lambda)}, \quad (3.10)$$

and the constants  $B_{\pm}$  and  $Q$  satisfy the relations

$$B_+ B_- = \frac{\kappa^2 Q^2 A(\lambda)}{(d-1)^2 (D-2)}. \quad (3.11)$$

We remind that we consider the case  $Q \neq 0$ . Due to eq. (3.3) we have

$$B_+ > B_- > 0. \quad (3.12)$$

In this case the  $(2+d)$ -dimensional section of the metric (3.6) has a horizon at  $r^{d-1} = B_+$ . For  $r^{d-1} = B_-$  the horizon is absent ( $\lambda \neq 0$ ). In the limit

$\lambda \rightarrow 0$  and  $D \rightarrow 2 + d$  we get the Myers-Perry  $O(d+1)$ -symmetric charged black hole solution [1]. For  $d = 2$ ,  $D = 4$  the solution coincides (up to redefinitions of the field variables) with the 4-dimensional dilatonic charged black hole solution [10,11].

## 4 Mass Bounds

The solution (3.6)-(3.11) describes an  $O(d+1)$ -symmetric charged dilatonic black hole with a chain of internal Ricci-flat spaces. The charge of the black hole is  $Q$  and the mass  $M$  is found from the time component of the metric (3.6) to be

$$2GM = B_+ + B_- \beta(\lambda), \quad (4.1)$$

where

$$\beta(\lambda) = 1 + 2\alpha_t = \frac{D-3-\lambda^2(D-2)}{D-3+\lambda^2(D-2)} \quad (4.2)$$

and

$$G = S_D \kappa^2 \quad (4.3)$$

is the effective gravitational constant ( $S_D$  is defined in [1]). It is clear that

$$-1 < \beta(\lambda) < 1 \quad (4.4)$$

and  $\beta(\lambda) = 0$  for  $\lambda_c^2 = (D-3)/(D-2)$ . Using the relation (4.1) and the inequalities (3.12) and (4.4) we get

$$M > M_c, \quad (4.5)$$

where

$$M_c = \frac{\kappa|Q|(D-3)}{G(d-1)\sqrt{A(\lambda)(D-2)}}. \quad (4.6)$$

This can be easily deduced from Fig. 1.

The critical case corresponds to

$$B_+ = B_- = \frac{\kappa|Q|\sqrt{A(\lambda)}}{(d-1)\sqrt{(D-2)}}. \quad (4.7)$$

Formula (4.6) agrees with the corresponding relation from ref. [14] in the case  $D = 4$ .

It is not difficult to verify that in the critical case (4.7) we have a horizon, if and only if

$$\lambda^2 \leq \frac{1}{d-1} - \frac{D-3}{D-2}. \quad (4.8)$$

For these values of  $\lambda$  the inequality (4.5) should be replaced by

$$M \geq M_c. \quad (4.9)$$

In the strong coupling limit  $\lambda^2 \rightarrow +\infty$  the critical mass (4.6) tends to zero. A possible interpretation of this effect seems to be screening by dilatonic field of the electric charge.

**Infinite-dimensional case.** Now we consider the interesting special case when  $n \rightarrow +\infty$ . In this limit the exact solution of the field equations taken from (3.6)-(3.11) reads

$$g = -f_+ f_-^{\frac{1-\lambda^2}{1+\lambda^2}} dt \otimes dt + f_-^{\frac{2}{d-1}} \left[ \frac{dr \otimes dr}{f_+ f_-} + r^2 d\Omega_d^2 \right] + \sum_{i=1}^n g^{(i)}, \quad (4.10)$$

$$F = Q r^{-d} dt \wedge dr, \quad (4.11)$$

$$\exp(2\lambda\varphi) = f_-^{\frac{1-\lambda^2}{1+\lambda^2}}, \quad (4.12)$$

where

$$B_+ B_- = \frac{\kappa^2 Q^2 (1 + \lambda^2)}{(d-1)^2}. \quad (4.13)$$

In this case the internal space scale-factors do not depend on the radial coordinate but the information about the presence of internal dimensions is

contained in the  $(2 + d)$ -dimensional part of the metric: this part does not coincide with the  $(D = 2 + d)$ -dimensional solution without internal spaces. The critical mass is non-zero in this limit:

$$M_c = \frac{\kappa|Q|}{G(d-1)\sqrt{1+\lambda^2}}. \quad (4.14)$$

## 5 Conclusion

We considered the  $O(d+1)$ -symmetric charged dilatonic black hole solution with  $n$  Ricci-flat internal spaces. We obtained the bound on the mass of a black hole for all (non-zero) values of the coupling constant. We found that the critical mass tends to zero in the strong coupling limit. We also considered the case of infinite number of internal spaces :  $n = \infty$ . In this case we obtained a non-trivial solution of the field equations and a non-zero value for the critical mass.

### Acknowledgments

The work was sponsored by the WIP project 016659/p. One of us (V. I.) was supported by DFG grant 436 RUS 113-7-2 and partly by the Russian Ministry of Science. V. I. also thanks the colleagues of the WIP gravitation project group at Potsdam University for their hospitality.



## References

- [1] R. C. Myers and M. J. Perry, Ann. of Phys., **172** (1986) 304.
- [2] U. Bleyer, K. A. Bronnikov, V. N. Melnikov and S. B. Fadeev, *On black hole stability in multidimensional gravity*, AIP preprint (Potsdam) 94-01, 1994.
- [3] V. D. Ivashchuk and V. N. Melnikov, *Multitemporal generalization of the Tangherlini solution*, submitted to Class. and Quantum Grav.; submitted to Izvestiya Vuzov (in Russian).
- [4] S. B. Fadeev, V. D. Ivashchuk and V. N. Melnikov, Chinese Phys.Lett., **8** (1991) 439; In: *Gravitation and Modern Cosmology*, Plenum Publ. NY, (1991) 37.
- [5] K. A. Bronnikov, Ann. der Phys., **48** (1991) 527.
- [6] K. A. Bronnikov, Izvestiya Vuzov, **No 1** (1992) 106 [in Russian].
- [7] S. B. Fadeev, V. D. Ivashchuk and V. N. Melnikov, *On Generalization of Charged Version of the Tangherlini Solution on the Case of  $n$  Internal Ricci-flat Spaces*, Nuclear Safety Institute preprint, Moscow, 1994; submitted to Izvestiya Vuzov (in Russian).
- [8] S. B. Fadeev, V. D. Ivashchuk and V. N. Melnikov, Phys. Lett., A **161** (1991) 98.
- [9] M. B. Green, J. H. Schwarz and E. Witten, *Superstring Theory*, Cambridge Univ. Press, 1986, p. 527.
- [10] G. W. Gibbons and K. Maeda, Nucl. Phys., **B298** (1988) 741.

- [11] D. Garfinkle, G. Horowitz and A. Strominger, Phys. Rev., **D43** (1991) 3140; **D45** (1992) 3888(E).
- [12] G. W. Gibbons and P. K. Townsend, *Vacuum Interpolation in Supergravity via Super  $p$ -branes*, DAMTP preprint R-93/19, 1993.
- [13] K. Shiraishi, Mod. Phys. Lett., **A 7** (1992) 3569.
- [14] G. W. Gibbons and C. G. Wells, *Anti-Gravity Bounds and the Ricci Tensor*, DAMTP preprint R93/25, 1993; to be published in Commun. Math. Phys.
- [15] N. S. Kalitzin, Wissenschaftliche Zeitschrift der Humboldt- Universitat zu Berlin, **Jg. VII** Nr 2 207 (1957/58).
- [16] V. D. Ivashchuk, Phys. Lett., **A 170** (1992) 16.

Figures:

a)

b)

Fig. 1: Lines of  $M = \text{const}$  for a)  $\beta > 0$  and b)  $\beta < 0$

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9405018v1>